

A Guide for Patients Beginning Buprenorphine Treatment at Home

Before you begin you want to feel very sick from your withdrawal symptoms

It should be at least . . .

- **12 hours** since you used heroin/fentanyl
- **12 hours** since snorted pain pills (Oxycontin)
- **16 hours** since you swallowed pain pills
- **48-72 hours** since you used methadone

You should feel at least three of these symptoms . . .

- Restlessness
- Heavy yawning
- Enlarged pupils
- Runny nose
- Body aches
- Tremors/twitching
- Chills or sweating
- Anxious or irritable
- Goose pimples
- Stomach cramps, nausea, vomiting or diarrhea

Once you are ready, follow these instructions to start the medication

DAY 1:

8-12mg of buprenorphine


Most people feel better the first day after 8-12mg. (Dosing depends on how early on the first day you started)

Step 1.

Take the first dose

Wait 45 minutes

4mg




- Put the tablet or strip under your tongue
- Keep it there until fully dissolved (about 15 min.)
- Do NOT eat or drink at this time
- Do NOT swallow the medicine

Step 2.

Still feel sick?
Take next dose

Wait 6 hours

4mg



Most people feel better after two doses = 8mg

Step 3.

Still uncomfortable?
Take last dose

Stop

4mg



- Stop after this dose
- Do not exceed 12mg on Day 1

DAY 2:

16mg of buprenorphine

Take one 16mg dose

Most people feel better with a 16mg dose

16mg

Repeat this dose until your next follow-up appointment

If you develop worsening symptoms while starting buprenorphine before your scheduled outpatient appointment return to the emergency department